arXiv:math/0612326v12 [math.MG] 9 Nov 2007

Steinhaus' problem on partition of a triangle

(to appear in Forum Geometricorum)

Apoloniusz Tyszka

Abstract. H. Steinhaus asked a question whether inside each acute triangle there is a point from which perpendiculars to the sides divide the triangle into three parts with equal areas. We present two methods of solving Steinhaus' problem.

The n-dimensional case of Theorem 1 was proved in [5], see also [2] and [3, Theorem 2.1, p. 152]. For an earlier mass-partition version of Theorem 1, for bounded convex masses in $\mathbb{R}^n$ and $r_1 = r_2 = ... = r_{n+1}$, see [6].

Theorem 1. Let $\boldsymbol{T} \subseteq \mathbb{R}^2$ be a bounded measurable set, and let $|\boldsymbol{T}|$ be the measure of $\boldsymbol{T}$. Let α_1, α_2, α_3 be the angles determined by three rays emanating from a point, and let $\measuredangle\alpha_1 < \pi$, $\measuredangle\alpha_2 < \pi$, $\measuredangle\alpha_3 < \pi$. Let r_1, r_2, r_3 be non-negative numbers such that $r_1 + r_2 + r_3 = |\boldsymbol{T}|$. Then there exists a translation $\lambda : \mathbb{R}^2 \to \mathbb{R}^2$ such that $|\lambda(\boldsymbol{T}) \cap \alpha_1| = r_1$, $|\lambda(\boldsymbol{T}) \cap \alpha_2| = r_2$, $|\lambda(\boldsymbol{T}) \cap \alpha_3| = r_3$.

H. Steinhaus asked a question ([9], [10]) whether inside each acute triangle there is a point from which perpendiculars to the sides divide the triangle into three parts with equal areas. Theorem 1 indicates only that there is such a point in the plane. There is no proof that partitions such as presented in Figure 1 and Figure 2 are not partitions into three parts with equal areas.



Figure 1



Figure 2

Long and elementary solutions of Steinhaus' problem appeared in: [7, pp. 101–104], [8, pp. 103–105], [11, pp. 133–138], [12]. For some acute triangle with rational coordinates of vertices, the point solving Steinhaus' problem is not constructible with ruler and compass alone, see [14]. Following article [13], we will present two methods of solving Steinhaus' problem. An anonymous referee found another two solutions which are constructive. His article will appear in this journal.

2000 Mathematics Subject Classification: 51M04, 51M25.

For $X \in \triangle ABC$ by $P(A,X)$, $P(B,X)$, $P(C,X)$ we denote figure areas at the vertices A, B, C respectively, see Figure 3. The areas $P(A,X)$, $P(B,X)$, $P(C,X)$ are continuous as functions of X on the triangle ABC. The function $f(X) = \min\{P(A,X), P(B,X), P(C,X)\}$ is also continuous. By Weierstrass' theorem f attains a supremum on the triangle ABC i.e. there exists $X_0 \in \triangle ABC$ such that $f(X) \leq f(X_0)$ for all $X \in \triangle ABC$.



Figure 3

Lemma. For a point X lying on a side of an acute triangle, the area at the opposite vertex is greater than some other area.

Proof. Without loss of generality, we may assume that $X \in \overline{AB}$ and $|AX| \leq |BX|$, see Figure 4. Straight line XX' parallel to straight line BC cuts the triangle AXX' greater than $P(A,X)$ (as the angle ACB is acute), but not greater than the triangle CXX' because $|AX'| < \frac{|AC|}{2} < |X'C|$. Hence $P(A,X) < |\triangle AXX'| \leq |\triangle CXX'| < P(C,X)$.



Figure 4

□

Theorem 2. If a triangle ABC is acute and f attains a supremum at X_0, then $P(A, X_0) = P(B, X_0) = P(C, X_0) = \frac{|\triangle ABC|}{3}$.

Proof. $f(A) = f(B) = f(C) = 0$, and 0 is not a supremum of f. Therefore X_0 is not a vertex of the triangle ABC. Let us assume, to set the attention, that $f(X_0) = P(A, X_0)$. By the Lemma $X_0 \notin \overline{BC}$. Suppose, on the contrary, that some of the other areas, let's say $P(C, X_0)$, is greater than $P(A, X_0)$.

Case 1: $X_0 \notin \overline{AC}$. When shifting X_0 from the segment $\overline{AB}$ by appropriately small ε and perpendicularly to the segment $\overline{AB}$ (see Figure 5), we receive $P(C, X)$ further greater than $f(X_0)$ and at the same time $P(A, X) > P(A, X_0)$ and $P(B, X) > P(B, X_0)$. Hence $f(X) > f(X_0)$, a contradiction.



Figure 5

Case 2: $X_0 \in \overline{AC} \setminus \{A, C\}$. By the Lemma

$$P(B, X_0) > \min\{P(A, X_0), P(C, X_0)\} \geq \min\{P(A, X_0), P(B, X_0), P(C, X_0)\} = f(X_0)$$

When shifting X_0 from the segment $\overline{AC}$ by appropriately small ε and perpendicularly to the segment $\overline{AC}$ (see Figure 6), we receive $P(B, X)$ further greater than $f(X_0)$ and at the same time $P(A, X) > P(A, X_0)$ and $P(C, X) > P(C, X_0)$. Hence $f(X) > f(X_0)$, a contradiction.



Figure 6

□

For each acute triangle ABC there is a unique $X_0 \in \triangle ABC$ such that $P(A, X_0) = P(B, X_0) = P(C, X_0) = \frac{|\triangle ABC|}{3}$. Indeed, if $X \neq X_0$ then X lies in some of the quadrangles determined by X_0. Let's say that X lies in the quadrangle with vertex A, see Figure 7. Then $P(A, X) < P(A, X_0) = \frac{|\triangle ABC|}{3}$.



Figure 7

The sets $R_A = \{X \in \triangle ABC : P(A, X) = f(X)\}$, $R_B = \{X \in \triangle ABC : P(B, X) = f(X)\}$ and $R_C = \{X \in \triangle ABC : P(C, X) = f(X)\}$ are closed and cover the triangle ABC. Assume that the triangle ABC is acute. By the Lemma $R_A \cap \overline{BC} = \emptyset$, $R_B \cap \overline{AC} = \emptyset$ and $R_C \cap \overline{AB} = \emptyset$. The theorem proved in [4] guarantees that $R_A \cap R_B \cap R_C \neq \emptyset$, see also [3, item D4, p. 101] and [1, item 2.23, p. 162]. Any point belonging to $R_A \cap R_B \cap R_C$ lies inside the triangle ABC and determines the partition of the triangle ABC into three parts with equal areas.

The above proof remains valid for all right triangles, because the thesis of the Lemma holds for all right triangles. For each triangle we have:

1) There is a unique point in the plane which determines the partition of the triangle into three equal areas.

2) The point of partition into three equal areas lies inside the triangle if and only if the thesis of the Lemma holds for the triangle.

3) The point of partition into three equal areas lies inside the triangle if and only if the supremum of f on the boundary of the triangle is smaller than the supremum of f on the whole triangle. For each acute or right triangle ABC, the supremum of f on the boundary does not exceed $\frac{|\triangle ABC|}{4}$.

4) The point of partition into three equal areas lies inside the triangle, if the triangle has two angles in the interval $(\mathrm{arctg}(\frac{1}{\sqrt{2}}), \frac{\pi}{2}]$. This condition holds for each acute or right triangle.

5) If the point of partition into three equal areas lies inside the triangle, then it is a partition into quadrangles.

For each triangle ABC with $\measuredangle C > \frac{\pi}{2}$ the point of partition into three equal areas lies inside the triangle if and only if

$$\sqrt{(1+\mathrm{tg}^2\measuredangle A)\cdot \mathrm{tg}\measuredangle B} + \sqrt{(1+\mathrm{tg}^2\measuredangle B)\cdot \mathrm{tg}\measuredangle A} > \sqrt{3\cdot(\mathrm{tg}\measuredangle A + \mathrm{tg}\measuredangle B)}$$

In case $\measuredangle C > \frac{\pi}{2}$ and

$$\sqrt{(1+\mathrm{tg}^2\measuredangle A)\cdot \mathrm{tg}\measuredangle B} + \sqrt{(1+\mathrm{tg}^2\measuredangle B)\cdot \mathrm{tg}\measuredangle A} = \sqrt{3\cdot(\mathrm{tg}\measuredangle A + \mathrm{tg}\measuredangle B)}$$

the unique $X_0 \in \overline{AB}$ such that $|AX_0| = \sqrt{\frac{(1+\mathrm{tg}^2\measuredangle A)\cdot\mathrm{tg}\measuredangle B}{3\cdot(\mathrm{tg}\measuredangle A+\mathrm{tg}\measuredangle B)}}\cdot|AB|$ and $|BX_0| = \sqrt{\frac{(1+\mathrm{tg}^2\measuredangle B)\cdot\mathrm{tg}\measuredangle A}{3\cdot(\mathrm{tg}\measuredangle A+\mathrm{tg}\measuredangle B)}}\cdot|AB|$ determines the partition of the triangle ABC into three equal areas. It is a partition into a triangle with vertex A, and a triangle with vertex B, and a quadrangle. There remains the case when $\measuredangle C > \frac{\pi}{2}$ and

$$\sqrt{(1+\mathrm{tg}^2\measuredangle A)\cdot \mathrm{tg}\measuredangle B} + \sqrt{(1+\mathrm{tg}^2\measuredangle B)\cdot \mathrm{tg}\measuredangle A} < \sqrt{3\cdot(\mathrm{tg}\measuredangle A + \mathrm{tg}\measuredangle B)} \qquad (*)$$

There is a straight line a perpendicular to the segment $\overline{AC}$ which cuts from the triangle ABC a figure with the area $\frac{|\triangle ABC|}{3}$, see Figure 8. There is a straight line b perpendicular to the segment $\overline{BC}$ which cuts from the triangle ABC a figure with the area $\frac{|\triangle ABC|}{3}$, see Figure 8. By $(*)$, the intersection point of straight lines a and b lies outside the triangle ABC. This point determines the partition of the triangle ABC into three equal areas.



Figure 8

References

[1] P. S. Alexandrov, *Combinatorial topology*, Dover Publications, Mineola, NY, 1998.

[2] K. Borsuk, *An application of the theorem on antipodes to the measure theory*, Bull. Acad. Polon. Sci. Cl. III 1 (1953), pp. 87–90.

[3] A. Granas and J. Dugundji, *Fixed point theory*, Springer-Verlag, New York, 2003.

[4] B. Knaster, K. Kuratowski and S. Mazurkiewicz, *Ein Beweis des Fixpunktsatzes fur n-dimensionale Simplexe*, Fund. Math. 14 (1929), pp. 132–137; reprinted in: K. Kuratowski, *Selected papers*, PWN (Polish Scientific Publishers), Warsaw, 1988, pp. 332–337, S. Mazurkiewicz, *Travaux de topologie et ses applications*, PWN (Éditions Scientifiques de Pologne), Warsaw, 1969, pp. 192–197.

[5] K. Kuratowski and H. Steinhaus, *Une application géométrique du théoréme de Brouwer sur les points invariants*, Bull. Acad. Polon. Sci. Cl. III 1 (1953), pp. 83–86; reprinted in: K. Kuratowski, *Selected papers*, PWN (Polish Scientific Publishers), Warsaw, 1988, pp. 520–523, H. Steinhaus, *Selected papers*, PWN (Polish Scientific Publishers), Warsaw, 1985, pp. 636–639.

[6] F. Levi, *Die Drittelungskurve*, Math. Z. 31 (1930), no. 1, pp. 339–345.

[7] E. Piegat, *Yet 105 problems of Hugo Steinhaus* (in Polish), Oficyna Wydawnicza GiS, Wrocław, 2000.

[8] E. Piegat, *Known and unknown problems of Hugo Steinhaus* (in Polish), Oficyna Wydawnicza GiS, Wrocław, 2005.

[9] H. Steinhaus, *Problem No. 132* (in Polish), Roczniki Polskiego Towarzystwa Matematycznego (Annales Societatis Mathematicae Polonae), Seria II, Wiadomości Matematyczne 9 (1966), no. 1, p. 99.

[10] H. Steinhaus, *Problem No. 779* (in Polish), Matematyka 19 (1966), no. 2, p. 92.

[11] H. Steinhaus, *Problems and reflections* (in Russian), Mir, Moscow, 1974.

[12] W. Stojda, *A solution of Problem No. 779* (in Polish), Matematyka 21 (1968), no. 5–6, pp. 267–273.

[13] A. Tyszka, *A solution of a problem of Steinhaus* (in Polish), Matematyka 49 (1996), no. 1, pp. 3–5.

[14] A. Tyszka, *Steinhaus' problem cannot be solved with ruler and compass alone* (in Polish), Matematyka 49 (1996), no. 4, pp. 238–240.

Apoloniusz Tyszka
Technical Faculty
Hugo Kołłątaj University
Balicka 116B, 30-149 Kraków, Poland
E-mail address: *rttyszka@cyf-kr.edu.pl*